



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

was observed to fly for several hundred yards, with characteristic speed and strength. That it was not a new comer is indicated by the fact that Mr. John Whitney, a man with a gunner's keen knowledge of the Anatinae, had reported some weeks earlier, a Pintail wintering in the vicinity. That the bird was strong of wing, after, evidently, surviving the low temperature, of the previous week, which ranged each night below zero Fahrenheit, indicates that it had remained through choice, rather than necessity, and with the foregoing notes, shows that the Pintail occasionally spends at least a considerable part of the winter as far northeast as Portland, Maine.—ARTHUR H. NORTON, *Portland, Me.*

White-winged Scoter (*Oidemia deglandi*) in Minnesota.—Late in the Autumn of 1905 a local hunter of this city shot an adult male White-winged Scoter on Lake Minnetonka. I examined this duck in the flesh and tried to buy it but the owner preferred to keep it and had it mounted by the late Henry W. Howling of Minneapolis. It was subsequently destroyed by a house cat.

On Nov. 14, 1911, I secured from a local hunter an adult female in perfect plumage, shot on Lake Minnewashta some two miles distant from this village. This was the only one seen and was not near any other species of duck. Cold weather with snow-storms had prevailed for three days prior to its capture. It was fat and in good condition.—ALBERT LANO, *Excelsior, Minn.*

The Bittern (*Botaurus lentiginosus*) Wintering in New Jersey.—On January 21, 1912, an adult Bittern was brought to me alive by my nephew, Robert W. Moore. It had been caught by him the day previous while he was skating at Walker's Pond about a mile and a half southeast of Haddonfield, N. J. For several days the bird had been noted about the pond in a weakened condition. No wounds were discoverable, but its stomach was empty and, I have no doubt, it was weak from lack of food. Two days after it was captured it died in captivity and was mounted.

Since the 5th of January snow had covered the ground and the extraordinary continuance of the freezing weather from that date to the 20th must have made the securing of its chosen food impossible. Possibly the unusual period of warm weather, which lasted throughout the early winter to January 5, tempted this bird, as well as many other species which usually go south, to winter with us. The latest previous record for the state seems to have been November 19, 1878, at a point much farther south in Cape May Co., and the earliest spring record March 30, so that this instance may fairly be termed the first recorded case of this species wintering in the state.—ROBERT THOMAS MOORE, *Haddonfield, N. J.*

Recent Occurrence of the Egret (*Herodias egretta*) near Portland, Maine.—Although a few authentic records have been made of the occurrence in Maine of the Egret (*Herodias egretta*), the increasing scarcity of